



**Case Report:**

**Idiopathic Infantile arterial calcification –A Very rare case**

Sharmila N, Professor,

Prashant. S. Joshi, Assistant Professor,

Dept. of Obstetrics and Gynecology, Adi Chunchangiri Institute of Medical Sciences, BG Nagara, Mandya, Karnataka, India,

Vani Ravikumar, Asst Professor Dept. of Pathology,

Bangalore Medical College, Bangalore

**Address For Correspondence:**

**Dr. Prashant. S. Joshi,**

Assistant Professor, Dept. of Obstetrics and Gynecology,

Adi Chunchangiri Institute Of Medical Sciences,

B.G. Nagara, Mandya, Karnataka, India.

**E-mail:** joshpj@rediffmail.com

**Citation:** Sarmila N, Joshi PS, Vani R. Idiopathic Infantile arterial calcification – A Very rare case. *Online J Health Allied Scs.* 2010;9(1):11

**URL:** <http://www.ojhas.org/issue33/2010-1-11.htm>

**Open Access Archives:** <http://cogprints.org/view/subjects/OJHAS.html> and <http://openmed.nic.in/view/subjects/ojhas.html>

Submitted: Mar 31, 2010; Accepted: Jun 10, 2010; Published: Jul 30, 2010

**Abstract:**

A rare case of Idiopathic Arterial Calcification of Infancy (IACI), inherited as an autosomal recessive disease, is reported.

**Key Words:** Arterial calcification, Infancy, Autosomal recessive

**Introduction:**


Calcification of heart and vessels in fetus is a very rare condition. It may be dystrophic or metastatic. An extremely rare form of vascular calcification has been termed Idiopathic Arterial Calcification of Infancy (IACI) which is inherited in an autosomal recessive pattern. We report a case of IACI diagnosed in our hospital.

**Case Report:**


A 26 year old woman, gravida 2, para 1 with no living child, with 20 weeks amenorrhea came for antenatal check up to our clinic. Her previous pregnancy was full term normal delivery and baby died at 11<sup>th</sup> month of age, reasons not being clear. Anomaly scan done at 20 weeks was reported as normal fetus. Patient came with preterm labor at 28 weeks of pregnancy and delivered a preterm baby weighing 1.2 kg which died 5 min after birth. Baby was sent for autopsy, as there was history of loss of first child. Autopsy report revealed fetal arterial calcification which involved massive calcification of aortic vessels, coronary vessels, pulmonary vessels, pancreatic vessels and vessels in the kidney. Liver showed extramedullary hematopoiesis and there was also hypertrophy of myocardium.




**Fig 1: Calcification of descending aorta**




**Fig 2: Calcification of aorta**



**Fig 3: Calcified pulmonary vessels**



**Fig 4: Calcified renal vessels**



**Fig 5: Myocardial hypertrophy**

are usually normal and there is no identifiable abnormality of calcium metabolism.<sup>2</sup>

**References:**

1. Saigal G. Idiopathic arterial calcification in infancy: pathology, diagnosis and treatment modalities. *Indian J Pediatr* 2002;69:265-267.
2. Stuart G. Idiopathic arterial calcification in two siblings: failure of treatment with diphosphonate. *Br Heart J* 1990;64:156-9.
3. Anderson KA, Burbach JA, Fenton LJ, Jaqua RA, Barlow JF. Idiopathic arterial calcification in infancy in newborn siblings with unusual light and electron microscopic manifestations. *Arch Pathol Lab Med* 1985;109:838-843.
4. Nagar AM. Antenatal detection of idiopathic arterial calcification with hydrops fetalis. *J Ultrasound Med* 2003;22:653-659.
5. Nasrallah MK. Prenatal diagnosis of idiopathic infantile arterial calcification with hydrops fetalis. *Ultrasound Obstet Gynecol* 2009;34: 601-604.

**Discussion:**

Idiopathic arterial calcification of infancy or occlusive infantile arterial calcification<sup>1</sup> is a rare congenital disorder which is usually fatal in first six months of life<sup>2</sup> and very few have survived more than one year. This condition is inherited as an autosomal recessive pattern<sup>3</sup> and diagnosis is usually made at autopsy in most of the cases.<sup>4</sup> Only about 161 cases have been described by literature to date, with antemortem diagnosis being made on the radiographic or sonographic demonstration of arterial calcification in only a few cases. It was first described by Bryant and White in 1901. It is characterized by widespread and extensive calcification and stenosis of large and medium sized arteries. This results from the deposition of calcium hydroxyapatite in the arterial internal elastic lamina layer, which leads to rapid progressive ischemic heart failure and refractory hypertension either prenatally or most commonly in neonatal and early infancy period.<sup>5</sup> Biochemical investigations